

Nonidentical fermions interact identically

Heather M. Hill

Citation: *Physics Today* **73**, 11, 14 (2020); doi: 10.1063/PT.3.4609

View online: <https://doi.org/10.1063/PT.3.4609>

View Table of Contents: <https://physicstoday.scitation.org/toc/pto/73/11>

Published by the *American Institute of Physics*

ARTICLES YOU MAY BE INTERESTED IN

[Magnetic field–boosted superconductivity](#)

Physics Today **73**, 44 (2020); <https://doi.org/10.1063/PT.3.4616>

[The math behind epidemics](#)

Physics Today **73**, 28 (2020); <https://doi.org/10.1063/PT.3.4614>

[Quantum computing ramps up in private sector](#)

Physics Today **73**, 22 (2020); <https://doi.org/10.1063/PT.3.4612>

[James Jeans and The Mysterious Universe](#)


Physics Today **73**, 36 (2020); <https://doi.org/10.1063/PT.3.4615>

[Atlantic invertebrates are going the wrong way](#)

Physics Today **73**, 17 (2020); <https://doi.org/10.1063/PT.3.4610>

[Does new physics lurk inside living matter?](#)

Physics Today **73**, 34 (2020); <https://doi.org/10.1063/PT.3.4546>



MCL
MAD CITY LABS INC.

Nanopositioning Systems Micropositioning AFM & SPM Single molecule imaging

The advertisement features a blue background with the MCL logo at the top center. Below the logo, four distinct pieces of precision engineering equipment are displayed in a row. From left to right: a large, light-colored square nanopositioning stage; a smaller, dark-colored micropositioning stage; a white AFM & SPM system with a striped calibration target; and a complex, dark-colored single molecule imaging setup with a camera and control unit.

Nonidentical fermions interact identically

The decoupling of electronic and nuclear spin states allows scattering fermionic atoms to rapidly cool.

Indistinguishable fermions don't typically interact much. Because their wavefunctions are antisymmetric, they tend to stay away from one another. That behavior manifests with particular strength in quantum gases cooled to ultralow temperatures, and it's a problem. Fermions' mutual avoidance limits the efficiency of evaporative cooling at low temperatures, as Deborah Jin observed 21 years ago when cooling potassium atoms (see *PHYSICS TODAY*, October 1999, page 17). Evaporative cooling relies on elastic collisions to nudge particles to lower energies, and it's an important technique for reaching the necessary temperatures to form Bose–Einstein condensates.

Bosons at ultralow temperatures can occupy the same energy state (see figure 1), which leads to physical proximity and much more scattering. What if fermions could act more like bosons? Not only would it be easier to evaporatively cool a Fermi gas down to the temperature needed for, say, quantum simulations, but there would also be new physics to explore. The challenge is finding fermions that don't mind sharing the same space.

Alkaline-earth fermionic atoms fit the bill. Because their nuclear spin states are decoupled from their electronic states, the en-

ergy levels and wavefunctions in an optical trap are identical for an atom in the $+\frac{1}{2}$ nuclear spin state, one in the $-\frac{1}{2}$ nuclear spin state, and so on. But the atoms aren't identical. As a result, fermionic atoms, each with a different nuclear spin value, can cluster together in the same energy state, as shown on the right in figure 1.

The interatom interactions are independent of the nuclear spin state, so for N atoms with different spins, the two-body interactions will have N -fold symmetry: The interaction is identical between any atom and each of the other $N-1$ atoms sharing its energy level. That so-called $SU(N)$ symmetry enhances those interactions and boosts otherwise weak effects into pronounced ones, with N as a tuning knob. Researchers are interested in $SU(N)$ -symmetric systems as a model to explore the physics behind a range of condensed-matter systems—for example, cuprates and other transition-metal oxides that display high-temperature superconductivity.

Research on $SU(N)$ Fermi gases started only in the past decade, and previous studies kept the gases in temperature regimes for which the interactions didn't significantly alter the system's thermodynamic behavior compared with non-

interacting Fermi gases that lack $SU(N)$ -symmetric interactions. Now the University of Colorado Boulder's Jun Ye, his graduate student Lindsay Sonderhouse, and their colleagues have cooled an $SU(N)$ Fermi gas down to a deeply degenerate regime, with all the states occupied below the Fermi level. Their ultracold strontium-87 gas with 10 distinct nuclear spin states shows clear signs of $SU(N)$ interactions and demonstrates a new way to rapidly cool fermions.¹

Clocking in

Ye and his colleagues have worked with ^{87}Sr gases for more than 16 years. Starting in 2014, they characterized the gas's scattering parameters and few-body elastic and inelastic interactions for clusters of up to five atoms.²

The Ye group's new study was motivated by three-dimensional optical lattice clocks, which were first demonstrated with a Fermi gas³ in 2017. (For more on optical-lattice clocks, see *PHYSICS TODAY*, March 2014, page 12.) 3D lattice clocks have the best intrinsic stability of any clock today. But getting a Fermi gas to the required low temperatures is a slow process. Speeding it up would improve clock performance because the detrimental influence of laser frequency noise depends on the time spent preparing the atoms.

The researchers predicted that the en-




FIGURE 1. BOSONS, FERMIONS, AND $SU(N)$ FERMIONS distribute themselves differently in energy levels at low temperatures. Whereas bosons can settle into the same state, fermions stick to separate levels below the Fermi energy E_F —unless they have interactions with $SU(N)$ symmetry. Those $SU(N)$ fermions can have N interacting particles in each level. (Adapted from ref. 1.)




FIGURE 2. LINDSAY SONDERHOUSE adjusts optical components for the laser-cooling experiment in Jun Ye's lab at the University of Colorado Boulder. (Photo by Christian Sanner.)

hanced scattering in an $SU(N)$ -symmetric Fermi gas would hasten cooling. So they timed it. To prepare their gas, the researchers used two stages of laser cooling to bring Sr atoms in a 2D potential down to $2 \mu\text{K}$. (A photo of the experimental setup is shown in figure 2.) They then introduced a single dimple trap for the atoms to pool into.

Their goal was to reach the highest density possible inside the dimple. Unfortunately, once atoms were in the dimple, the cooling light created collisions and effective repulsion between them. To overcome that problem, Ye and his colleagues borrowed a technique used for bosons:⁴ They added another laser that rendered the atoms inside the dimple transparent to the cooling laser. The transparency laser shifted the excited state of the atoms such that the cooling light was no longer resonant with the transition to the ground state. Atoms outside the dimple continued to be optically cooled, and the ones inside thermalized with them.

The researchers then left the $SU(N)$ -symmetric Fermi gas to evaporate for either 600 ms or 2.4 s, depending on the desired temperature. That evaporation time is down from about 10 s for two-spin gases. They discovered that the more fermions that are colliding in each energy level in the dimple trap, the faster they cool; the total evaporation time scales

approximately as $1/(N-1)$. The researchers prepared atoms with all 10 possible nuclear spin states—that is, 10 atoms per energy level—and 5×10^4 atoms per spin state by the end of a 2.4 s evaporation.

Dynamic changes

Although the fermion interactions are weak, they measurably change the system's thermodynamics. One example is the gas's compressibility. The $SU(N)$ interactions are repulsive in ^{87}Sr . And although the compressibility depends only weakly on the number of atoms per spin state, it scales as $N-1$ with the number of spin states.

Ye and his group measured the gas's compressibility from its density fluctuations. To do so, they released the gas from the trap and performed absorption imaging, which gives a snapshot of the density profile. They found that the compressibility was reduced by 18% for $N=10$ compared with the model for a noninteracting Fermi gas at the same density and temperature.

But the density profile alone can't confirm that the system's constituents interact. The fluctuations for an interacting system look the same as a colder noninteracting system, although the mechanisms that limit compression are different: interaction-induced repulsion in the one case and filled energy levels in the other.

TENURE-TRACK FACULTY POSITIONS IN EXPERIMENTAL AND THEORETICAL PHYSICS

The Department of Physics invites applications for tenure-track faculty positions at the Assistant Professor level. Ranks at Associate Professor or above will also be considered for candidates with exceptional record of research excellence and academic leadership. Applicants must possess a PhD degree in physics or related fields and have evidence of strong research productivity.

We seek experimental candidates in quantum matter and quantum information, including quantum and low-dimensional materials, materials with strong electronic correlations, cold atoms, quantum optics, and quantum enabled technologies. We also seek theoretical candidates in condensed matter theory, statistical physics, neural networks or data analytics.

Appointees are expected to assume teaching responsibilities for undergraduate and graduate courses, and to conduct vigorous research programs. Further information about the Department is available at <http://physics.ust.hk>.

Starting salary will be commensurate with qualifications and experience. Fringe benefits including medical and dental benefits, annual leave and housing will be provided where applicable. Initial appointment will normally be on a three-year contract. A gratuity will be payable upon successful completion of contract.

Application Procedure

Applicants should submit their application including CV, cover letter, complete publication list, research statement, teaching statement, and three reference letters, via AcademicJobsOnline.Org at (<https://academicjobsonline.org/ajo/jobs/16290>).

Please quote reference number "PHYS2509" in your application materials.

Screening of applications begins immediately, and will continue until the positions are filled.

The University of Texas at Austin invites applications for multiple tenure-track Assistant Professor positions (subject to the availability of funding) to enhance its research programs in the general area of quantum materials and quantum information science. The candidates will have a home department in Physics or Computer Science. This hiring initiative focuses on two areas: (1) fundamentals and applications of quantum information, algorithms, systems and architectures, and (2) quantum materials, devices, and quantum simulation.

The positions offer excellent start-up funds, salary, and laboratory space in a dynamic and highly interactive research environment.

Application Instructions:

Please submit your application at apply.interfolio.com/79208 and include: a cover letter, a description of research accomplishments and current and future research plans (limited to 4 pages), a statement on teaching experience, goals, and philosophy, a Curriculum Vitae, and a list of publications. Candidates should indicate a preferred home department and are also required to request three letters of recommendation using the Interfolio website, which will email the referees with instructions to directly upload their letters of recommendation. Complete applications will be reviewed on an on-going basis beginning September 30, 2020.

Applications received before November 30 will be given full consideration.

The University of Texas at Austin, as an equal opportunity/affirmative action employer, complies with all applicable federal and state laws regarding nondiscrimination and affirmative action. The University is committed to a policy of equal opportunity for all persons and does not discriminate on the basis of race, color, national origin, age, marital status, sex, sexual orientation, gender identity, gender expression, disability, religion, or veteran status in employment, educational programs and activities, and admissions.




FIGURE 3. SIGNATURES OF FERMION INTERACTIONS in a strontium gas cloud show up as anisotropies in experimental data and theory. After the interacting Fermi gas is released from an optical trap, absorption images record a snapshot of the density profile that may appear anisotropic. But plotting the density profiles' anisotropy reveals a lobed structure in the experimental data and the interacting theory. (Adapted from ref. 1.)

Verifying the presence of interactions requires additional information, such as the expansion dynamics. When an ideal, noninteracting Fermi gas is released from a trap, even an asymmetric one, the gas's profile ends up isotropic because the isotropic momentum distribution produces uniform ballistic expansion. Interactions change the gas's kinetic energy and preferentially propel the atoms along the direction with the largest density gradient, arising from the asymmetry in the trap. As a result, an interacting Fermi gas has a nonuniform distribution. The researchers' gas of ^{87}Sr atoms with 10 different nuclear spin states yielded a noticeably elliptical cloud despite the weakness of the interactions.

Gases with an isotropic density distribution appear circular long after they're released from the trap. But the expansion dynamics of noninteracting and interacting gases still subtly differ. The team's kinetic theory calculations, performed by Ana Maria Rey, show that the density profile n of each gas appears to be circularly symmetric and therefore not indicative of interactions, or the lack thereof. But the differences become clear from the distributions' transpose anisotropy, defined as $n(x, z) - n(z, x)$, shown in figure 3.

In the interacting model and the experimental data, clear lobes emerge in the anisotropy that are not present in the noninteracting model. And because the lobes decrease with increasing temperature, integrals over the transpose anisotropy can be used as a temperature probe with an accuracy of $\frac{1}{100}$ of the Fermi temperature—the temperature at which thermal effects are comparable to quantum effects, defined as $T_F = E_F/k_B$ in terms of the Boltzmann constant and the Fermi

energy, or the energy difference between the highest and lowest states occupied by noninteracting fermions at absolute zero temperature.

A new spin

With an efficient preparation method and a basic understanding of their properties, researchers are ready to tackle problems in condensed-matter and high-energy physical systems with $SU(N)$ -symmetric gases. "One can almost think of a deeply degenerate Fermi gas as premium fuel for a quantum simulator," says Ye. Different combinations of kinetic energy, interaction energy, and nuclear spins can be used to systematically explore the phase diagram of, for example, the Fermi-Hubbard model (see the article by Gabriel Kotliar and Dieter Vollhardt, *PHYSICS TODAY*, March 2004, page 53), whose phase diagram is still unknown despite its common use to describe strongly correlated materials.

The system in the current study follows the Fermi-liquid model for interacting fermions, which describes the normal state of most metals at low temperatures. But in the future, $SU(N)$ -symmetric gases may move to regimes where the theory isn't valid. Exploring the physics in that regime may help researchers understand the origins of the quantum phase transition in heavy-fermion materials.

Heather M. Hill

References

1. L. Sonderhouse et al., *Nat. Phys.* (2020), doi:10.1038/s41567-020-0986-6.
2. X. Zhang et al., *Science* **345**, 1467 (2014); A. Goban et al., *Nature* **563**, 369 (2018).
3. S. L. Campbell et al., *Science* **358**, 90 (2017).
4. S. Stellmer et al., *Phys. Rev. Lett.* **110**, 263003 (2013).